

CONTENTS

Introductory lectures	pages
1. Falcinelli M, Provorov N, Tikhonovich I, Veronesi F. Sustainable crop production: scientific roots and prospects	11
2. Romanenko GA. Impact of Russian Academy of Agricultural Sciences in cultivation of fodder crops and in fodder production in Russia	14
Section 1. Germplasm Evaluation and Utilization	
3. Simon U. Indigenous grassland as a genetic resource for breeding forage plants	21
4. Zhuchenko AA. Ecological and genetic foundations for adaptive system of breeding fodder crops	24
5. Bullitta S, Meloni MC, Piluzza G. Alternative pasture legumes from Asinara Island (North Western Sardinia)	38
6. Posselt UK, Bolaric S. Genetic diversity among cultivars and ecotypes of <i>Lolium perenne</i> L.	40
7. Wilkins PW. Effect of extreme leafiness introgressed into a forage-type perennial ryegrass variety on herbage quality and yield	41
8. Campbell MC, Francis CM, Kailis SG, Clements J, Milton JTB. Some exotic oilseeds with fodder potential	44
9. Dzyubenko NI, Malyshev LL. The taxonomic and ecological diversity of amenity grasses in Russia and other former Soviet Union states	46
10. Kolomeichenko VV. Use of information about the productive process in breeding sustainable fodder crops and amenity grasses	47
11. Vishnyakova M. Grain legumes germplasm collection of Vavilov Institute as a source for breeding of adaptive varieties of fodder crops	49
12. Pacucci G, Troccoli C, Leoni B. Evaluation of new common vetch (<i>Vicia sativa</i> L.) genotypes for forage yield in Southern Italy. 1 - Influence of seed size and planting depth	50
13. Pacucci G, Troccoli C. Evaluation of new common vetch (<i>Vicia sativa</i> L.) genotypes for forage yield in Southern Italy. 2 - Effect of plant density	52
14. Alexandrova TG. Intraspecific diversity of <i>Vicia angustifolia</i> Reichard and <i>Vicia cordata</i> Wulf. ex Hoppe - the closest wild-growing relatives of common vetch (<i>Vicia sativa</i> L.)	54
15. Alexandrova TG, Potokina EK, Francis CM, Bennett SJ. Genetic database of common vetch (<i>Vicia sativa</i> L.) from the area of Former Soviet Union in the collection of N.I.Vavilov Institute of Plant Industry	55
16. Turin YS, Fitsev AI, Michaylichenko AB. Main directions of the vetch breeding	56

17. Yan'kov I, Andreeva N. Agronomic characters of <i>Vicia ervilia</i> germplasm collection maintained in the Vavilov Institute of Plant Industry	58
18. Voronichev BA, Kolomeichenko VV. Main trends and breeding methods of field beans (<i>Vicia faba</i> L.) in Russian Federation	59
19. Burlyaeva MO. Selection of the most informative parameters for evaluation of odder soybean by green biomass productivity	60
20. Burlyaeva MO, Malyshev LL. The informative characters for breeding soybeans of different fodder usage	62
21. Burlyaeva MO, Chmeleva ZV. The biochemical value of green biomass of grass pea (<i>Lathyrus sativus</i> L.) introduced in Leningrad Region	64
22. Staszewski Z, Bodzon Z. Inheritance of branch raceme inflorescence determined by a spontaneous mutation in <i>Medicago sativa</i> L.	65
23. Bender A, Tamm S. Polymorphism for fertility and self-fertility in alfalfa varieties Jogeva 118 and Ellerslie I	68
24. De Vliegher A, Carlier L, Van Waes C, Van Waes J. Varietal differences in yield and nitrogen efficiency in <i>Lolium perenne</i>	70
25. Lengliz W, Chakroun M, Zouaghi M, Rezgui S. Agronomic evaluation of introduced oat lines	73
26. Abraham HM. The efficiency of mass honeycomb selection for yield in <i>Dactylis glomerata</i> L. and <i>Agropyron cristatum</i> L.	76
27. Annicchiarico P, Lucaroni B, Piano E, Russi L, Veronesi F. An Italian network for evaluation of the turf species and varieties	78
28. Funtova VG. Grassland ecosystems as a source of perspective fodder crop populations for breeding	81
29. Pelikan J, Zapletalova I, Hofbauer J, Nedelnik J, Pokorny R. Study of fodder plant collections in the Research Institute for Fodder Plants Ltd., Troubsko, Czech Republic, in 1994-1999	82
30. Odoardi M, Paoletti R. Forage potential, persistence and quality factors of alpine populations and varieties grown over the timber line	84

Section 2. Adaptive Potential and Environmental Manipulations

31. Michaylichenko BP. Ecological and evolutionary approaches and adaptive methods for breeding fodder crops in ecologically based agriculture in Russia	89
32. Baert J, Lamote V, Van Bockstaele E. Searching for meiotic tetraploid perennial ryegrass	91
33. Shamsutdinov ZS. Ecotypical plant breeding concept for creation of ecologically differentiated varieties of xerophytes and halophytes	94
34. Shevelukha VS. Biological objects as self-regulating systems	96

35. Piskovatsky YM, Stepanova GV. Modern methods in phytocenotic and symbiotic breeding of alfalfa	98
36. Breazu IT, Oprea G, Balan M, Neagu M. The use of allelopathy to sustain development in herbage legumes breeding	100
37. Novoselov M, Drobysheva L, Zyatchina G. Modern methods in breeding of red clover: reproduction, symbiotic and soil aspects	104
38. Nagy B. The genetical possibilities of the increasing of persistence in alfalfa (<i>Medicago sativa</i> L.)	105
39. Gau M, Sawai A, Uchiyama K, Yamaguchi H. Combinig ability in forage yield in alfalfa (<i>Medicago sativa</i> L.) from ecotypic population in Northern part of Japan	107
40. Tumasova M, Krylova L, Beresneva O. Breeding of red clover for tolerance against environmental stress factors of north-east European region of Russia	109
41. Pozdnyakov VA, Drizhachenko AI. Impact of environmental factors on induction of flowering in the interspecific hybrids <i>Trifolium pratense</i> L. x <i>T. diffusum</i> Ehrh.	110
42. Ghesquiere M, Marhadour S, Mousset Cl. Through selection against flowering in introgressed ryegrass x fescue progenies	113
43. Marhadour S, Ghesquiere M. Rooting aspects in persistency of ryegrass x fesque hybrids	115
44. Tsvetova MI. Characteristics of sorghum pollen from the point of view of estimation of diverse influences upon plant	117
45. Klimont K, Rafalski A, Wisniewska I. The influence of laser biostimulation on the yield and seed quality of amaranth (<i>Amaranthus cruentus</i> L.)	118
46. Nossov N. Chemical and agrotechnical approaches to decrease content of dandelion (<i>Taraxacum officinale</i>) in long-term grasslands	122

Section 3. Benefits from Microbes

47. Lugtenberg BJJ. Use of bacteria for plant growth promotion and plant protection	127
48. Tikhonovich IA, Kozhemyakov AP, Ovtyna AO, Provorov NA. Construction of highly efficient symbiotic systems	131
49. Minchin FR. The regulation of nitrogen fixation in legumes	136
50. Provorov NA. Co-ordinated plant-microbe breeding: a key approach for making legume crops sustainable	139
51. Onishchuk OP, Kurchak ON, Sharypova LA, Krol EA, Zatovskaya TV, Simarov BV. Genetics of competitiveness of <i>Sinorhizobium meliloti</i>	141

52. Khodjaiova L, Ramenskaya M, XiChun, Lutova L. A new two-step procedure for selection of plants resistant to the sterol-dependent pests and pathogens	144
53. Belimov AA, Kunakova AM, Safranova VI, Kozhemyakov AP, Yudkin LV, Stepanok VV, Khamova OF, Postavskaya SM. Interaction between associative bacteria and barley under environmental stresses: input of partners' genotypes and growth conditions	146
54. Metlitskaja J, Provorov NA, Meripold H, Rausberg P, Laitamm H. Selection of goat's rue (<i>Galega orientalis</i> Lam.) for increasing efficiency of symbiosis	148
55. Kurchak ON, Galan MS, Orlova OV, Provorov NA, Simarov BV. A potential for symbiotic nitrogen fixation in hairy vetch (<i>Vicia villosa</i> Roth) as compared with common vetch (<i>Vicia sativa</i> L.)	151
56. Labutova NM, Malinovsky BN, Poyda VB, Smilovenko LA. Impact of wheat (<i>Triticum monococcum</i> , <i>Triticum</i> spp.) and sorghum (<i>Sorghum vulgare</i>) genotypes on the efficiency of symbiosis with <i>Glomus intraradices</i>	155
57. Govedarica M, Jarak M, Milosevic N. Effectiveness of <i>Rhizobium meliloti</i> in alfalfa: nodulation and microbial activity of soil	157
58. Milosevic N, Jarak M, Govedarica M. Field testing of <i>Rhizobium leguminosarum</i> bv. <i>viceae</i> for effectiveness in forage peas	159
59. Maslenikov SE. Increase of alfalfa nitrogen-fixing apparatus as a result of influence of fungi - root rot pathogens	161
60. Tolkachev NZ, Sichkar VI. Intracultivar soybean selection for improvement of nitrogen fixation	162
61. Chernyaeva II, Tuev NA, Vasilieva ND, Kunakova AM, Kozhemyakov AP. The interaction of alfalfa and clover with <i>Rhizobium</i> on different mineral nitrogen backgrounds	163
62. Kravchenko LV, Strigul NS, Azarova TS, Tikhonovich IA. Experimental and mathematical models of the microbial-plant system in creating new criteria for plant breeding	165
63. Patyka VF, Nadkernichnaya EV, Tolkachev NZ. Nitrogen fixing potential of winter rye and soybean and its using in selection	168
64. Kasakova AS. Effect of vesicular-arbuscular mycorrhizal inoculation on growth of sorghum plants	169
65. Sviridova OV, Vorobyov NI, Pishchik VN, Mikhaleva LV. Effect of humic preparations on plant-microbial associations	171
66. Takai T. Comparison between heat treated and non-treated seeds for freezing tolerance and snow mold resistance in meadow fescue (<i>Festuca pratensis</i> Huds.)	173

Section 4. Genome Analysis and Modification

67. Lorenzetti F. Genome analysis and modification (Section 4 Chairpersons' remarks)	177
68. Falcinelli M, Barcaccia G. Assisted-selection for apomixis in <i>Poa pratensis</i> L. by PCR-based molecular markers	179
69. Rognli OA, Alm V, Busso C. Construction of an AFLP- and RFLP-based linkage map in meadow fesque (<i>Festuca pratensis</i> Huds.)	183
70. Barcaccia G, Albertini E, Veronesi F. Inter-microsatellite DNA markers in the <i>Medicago sativa</i> complex	185
71. Agafodorova MN, Solodkaya LA, Koreneva IV, Kopertekh LG, Ivashuta SI. The using of genetic engineering methods for fodder crop breeding	191
72. Abberton MT. Molecular marker analysis in white clover (<i>Trifolium repens</i> L.)	192
73. Humphreys MO. The use of marker assisted selection in ryegrass breeding	196
74. Thorogood D, Paget M, Humphreys M, Turner L, Roderick H. QTL analysis of crown rust resistance in perennial ryegrass	200
75. Rosellini D, Veronesi F, Bingham ET. Male fertility assessment within and between lucerne cultivars by pollen racing	207
76. Isobe S, Gau M, Kozlov N, Ivashuta S, Agafodorova M, Uchiyama K. Construction of red clover genetic linkage map using RFLP markers	210
77. Kolliker R, Jones ES, Drayton M, Dupal M, Forster JW. Development and implementation of AFLP and SSRP markers in white clover	211
78. Elkonin LA. Genetic analysis and use of new sources of male sterility induced in tissue culture of sorghum	213
79. Rognli OA, Nurminniemi M, Nilsson N-O, Tufto J. Pollen dispersal and gene flow in grasses – a model study related to the risk of gene flow from the use of future transgenic grasses	215
80. Alm V, Busso C, Rognli OA. Linkage mapping in an outbreeding full-sib family of <i>Festuca pratensis</i> Huds.: theory and application	217
Attached Workshop «<i>Pisum sativum</i> - a Tool for Studying Plant-Microbe Interactions»	
81. Priefer UB. Basic mechanisms in the interaction between micro- and macrosymbiont in the <i>Rhizobium leguminosarum</i> - pea symbiosis: the bacterial side	221
82. Lie TA, Hartati Y. The use of mini and twin plants for the genetic analysis of symbiotic variation of <i>Pisum sativum</i> L. towards <i>Rhizobium leguminosarum</i> strains	223
83. Tempelman-Bobbink A, Yami KD, Lie TA. Symbiotic diversity of <i>Pisum sativum</i> L. ecotype <i>humile</i> and the associated <i>Rhizobium leguminosarum</i> strains	225

84. Provorov NA, Tikhonovich IA, Kozhemyakov AP. Co-ordinated breeding of <i>Pisum sativum</i> plants and <i>Rhizobium leguminosarum</i> strains for the effective symbiosis	228
85. Borisov AY, Jacobi LM, Lebsky VK, Morzhina EV, Tsyganov VE, Voroshilova VA, Tikhonovich IA. Pea (<i>Pisum sativum</i> L.) genetic system controlling development of nitrogen-fixing nodules and arbuscular mycorrhiza	231
86. Jacobi LM, Kukalev AS, Ushakov KV, Tsyganov VE, Provorov NA, Borisov AY, Tikhonovich IA. Genetic variability of garden pea (<i>Pisum sativum</i> L.) for symbiotic capacities	237
87. Thomas CL, Harrison S, Jones A-L, Revers F, Maule AJ. Exploiting host: virus interactions using pea seed borne mosaic virus (PSbMV) and pea	240
88. Francis C, Khan T, Priliouk L, Gorfu D, Bejiga G. Screening peas of Vavilov Institute and ICARDA for black spot resistance	242
89. Ovtysna AO, Tikhonovich IA, Staehelin C, Schultze M, Kondorosi A. Rhizobial Nod factors induce their own degradation by pea roots	244
90. Lutova LA, Pavlova ZB. Role of phytohormones in pea-rhizobia and pea-agrobacteria interactions	246
91. Belimov AA, Safranova VI, Tsyganov VE, Borisov AY, Soboleva VN, Kvokova NA, Zaitseva LN, Poddubnykh ON, Stepanok VV, Dolgaya LN, Sukhanov PA, Tikhonovich IA. Screening of garden pea varieties by resistance to cadmium and accumulation of heavy metals	248
92. Rozov SM, Borisov AY, Tsyganov V, Kosterin O. The history of the pea gene map: last revolutions and the new symbiotic genes	251
93. Naumkina T, Yakovlev V, Titonok T, Vasilchikov A, Orlov V, Borisov A, Kulikova O. Pea breeding to improve effectiveness of symbiotic nitrogen fixation	254
94. Serdyuk V. New sources of germplasm for the fodder pea (<i>Pisum sativum</i> L.) breeding	256
95. Priliouk L, Robertson L, Sweid F. Evaluation of Vavilov Institute pea collection for cold tolerance in Syria	258
96. Krugova ED, Mandrovskaya NM, Okhrimenko SM. Biologically active substances for increasing the efficiency of pea nitrogen fixation	259
97. Borisov AY, Boesten B, Sanjuan J, Sanjuan-Pinilla JM, Tsyganov VE, Voroshilova VA, Olivares J, Priefer UB, Tikhonovich IA. Pea (<i>Pisum sativum</i> L.) symbiotic gene mutations affecting the expression of <i>Rhizobium leguminosarum</i> bv. <i>viceae</i> symbiotic genes	259
Index of Latin names	267
Author index	273
List of participants	276